

Plant Growth-Rate Dependence of Detrital Carbon Storage in Ecosystems

Just Cebrián* and Carlos M. Duarte

Plant Growth-Rate Dependence of Detrital Carbon Storage in Ecosystems

Just Cebrián* and Carlos M. Duarte

Detrital carbon accumulation accounts for most of an ecosystem's capacity to store organic carbon because the carbon contained as plant detritus exceeds that stored in living plants by about threefold. A comparative analysis of the mass and turnover of detrital carbon in ecosystems demonstrates that these properties are strongly related to the turnover rate of the dominant primary producers and are poorly related to ecosystem primary production. These results contribute to an understanding of the factors that control carbon storage in ecosystems and the role of carbon storage in the global carbon budget.

The assessment of factors that control C storage in ecosystems (1-3) is essential for determining the role of vegetation in the global C budget (4, 5). Carbon storage in ecosystems is accounted for mostly by the detrital C mass, which amounts to about threefold that accounted for in living plant tissues (3, 4). Hence, knowledge of the factors that control the size and turnover of the detrital C pool in ecosystems should help elucidate the processes that control C sinks in the global C budget (5).

In this report we use a broad-scale comparison from published values to show that even though detrital C flux is strongly controlled by ecosystem primary production, neither one is strongly related to the mass and turnover of the detrital C pool. We then demonstrate that plant turnover rate explains a major fraction of the variance in detrital C mass and turnover among ecosystems. We compiled data from reports on aboveground biomass and primary production and the mass and production of detrital C (6) from a broad range of ecosystems (7-11).

Carbon flux into the detrital pool was strongly and linearly related to primary pro-

duction (Fig. 1). This linear relation applied both to individual ecosystem types and to all the types grouped together (12). On average ~56% of primary production enters the detrital pool, and, with the exception of grasslands, more productive ecosystems yield a correspondingly higher C flow into the detrital pool. Our results support the general findings that litterfall rates are higher in more productive forests (13) and that phytoplanktonic primary production is positively correlated to phytoplankton sedimentation rates (14).

Detrital C mass was poorly related to both primary production and C flux into the detrital pool (Fig. 2), both of them explaining only ~10% of the variation in detrital C mass among ecosystems. The C mass of the detrital pool varied by about three orders of magnitude for similar values of C flux into this pool (Fig. 2B). The differences in detrital C mass among ecosystems were instead strongly correlated to the plant turnover rate (Fig. 3A); the tendency toward reduced detrital C mass with increasing plant growth rate accounted for 53% of the variance in detrital C mass.

The observations that detrital C mass is relatively independent of C flux into the detrital pool (Fig. 2B), whereas it declines as plant growth rate increases (Fig. 3A) imply that the loss rate of detrital C

Centro de Estudios Avanzados de Blanes, Consejo Superior de Investigaciones Científicas, Camí de Santa Bàrbara s/n, 17300 Blanes, Girona, Spain.

*To whom correspondence should be addressed.




Fig. 1. The relation between C flux into the detrital pool and primary production (both in units of grams of C per square meter per day). The solid line depicts the regression equation fitted to the data. Symbols represent (*) plankton, (●) benthic microalgae, (Δ) macroalgal beds, (▲) freshwater macrophyte meadows, (■) seagrass meadows, (□) grasslands, and (○) forests.

should increase with increasing plant growth rate. The net result of these processes is a significant tendency toward an increase in the turnover rate of the detrital C pool with increasing plant growth rate (Fig. 3B). This finding is consistent with the fast decomposition rates reported for the nutrient-rich detritus produced by fast-growing plants (15). Export of detrital C tends to be greater in aquatic ecosystems than in terrestrial ecosystems, but it typically represents only a small fraction of the C loss from the detrital pool (9, 10); hence, export of detrital C cannot have a significant effect on the observed patterns.

Our results indicate that plant turnover rate controls the size and turnover of the detrital C pool, thereby setting the capacity of an ecosystem to store C. Hence, ecosystems dominated by slow-growing plants accumulate large, slowly decomposing detrital pools which act as C sinks both on a local and global scale (1, 2). In contrast, C accumulation in the detrital pool of ecosystems dominated by fast-growing plants is much smaller. The magnitudes of primary production and the flux of C entering the detrital compartment reveal little about an ecosystem's capacity to store detrital C (Fig. 2).

The relation between detrital C mass and plant turnover rate described here provides a basis to assess the response of the detrital C pool to global changes in plant turnover rate. Several lines of evidence suggest that global changes in land use and climate may be conducive to a global increase in plant turnover rate (1, 3, 16–20). Coastal eutrophication has been shown to cause a shift from slow-growing, thick macroalgae and seagrasses to fast-growing phytoplankton and mac-




Fig. 2. The relation between detrital C mass (grams of C per square meter) and (A) primary production (grams of C per square meter per day), and (B) the C flux into the detrital pool (grams of C per square meter per day). The detrital C mass is poorly, although significantly ($P < 0.05$), correlated to the primary production ($r^2 = 0.1$, where r is the Pearson correlation coefficient) and to the C flux into the detrital pool ($r^2 = 0.13$). Symbols as in Fig. 1.

roalgae (16). Deforestation replaces slow-growing plants (that is, forests) by faster growing (that is, grasslands and agricultural crops) plants (1, 3, 17). Grasslands are expected to expand through the northeastern American territories, which are presently colonized by mixed conifer-hardwood forests; in response to an atmospheric warming of 1.5° to 4.5°C by the end of the century (18). The turnover rate of tropical forests has increased since the 1950s (19), and higher atmospheric CO₂ concentrations may lead to a further increase in plant turnover rate (20). Our results suggest that these trends toward a global increase in plant turnover rate could result in a net decline in ecosystem C storage from losses of soil C. This prediction is in agreement with results from simulation models that combine the effects of change in climate, atmospheric composition, and the global spread of agricultural crops and range lands (1).

Our prediction of the effect of global changes in plant turnover rate on the size of the detrital C pool remains, however, qualitative because our data do not represent a random sample of the different ecosystems on Earth. Furthermore, the patterns de-




Fig. 3. (A) The relation between detrital C mass (C_{detrital} , grams of C per square meter) and plant turnover rate [PP/B , per day, where PP is primary production (grams of C per square meter per day) and B is plant biomass (grams of C)] for different ecosystems. The solid line represents the regression equation fitted to the data: $\log C_{\text{detrital}} = (0.43 \pm 0.24) - (0.67 \pm 0.08) \log PP/B$ (with $r^2 = 0.53$, $n = 53$, F test, $P < 0.00001$). (B) The relation between detrital C turnover rate (TC_{detrital} , per day) and plant turnover rate for different ecosystems. The solid line represents the regression equation fitted to the data: $\log TC_{\text{detrital}} = (-0.9 \pm 0.2) + (0.64 \pm 0.08) \log PP/B$ (with $r^2 = 0.52$, $n = 53$, F test, $P < 0.00001$). Symbols as in Fig. 1.

scribed here represent empirical relations that may be blurred by feedback effects associated with simultaneous changes in growth conditions in response to changing global temperature and atmospheric CO₂ concentration (21). Yet, we demonstrate that the size and turnover rate of the detrital C pool are closely related to the turnover rate of the plant community and are poorly related to ecosystem primary production. Thus, the mechanisms linking the dynamics of the detrital C pool to plant turnover rate must be investigated to improve our capacity to model C storage in ecosystems and to evaluate its role in the global C budget.

REFERENCES AND NOTES

1. R. Dixon et al., *Science* 263, 185 (1994).
2. S. Wofsy et al., *ibid.* 260, 1314 (1993); R. Birdsey, A. Plantinga, L. Heath, *For. Ecol. Manage.* 58, 33 (1993).
3. H. Eswaran, E. Van Den Berg, P. Reich, *Soil Sci. Soc. Am. J.* 57, 192 (1993).
4. E. Sundquist, *Science* 259, 934 (1993); U. Slen-

- genthaler and J. L. Sarmiento, *Nature* 365, 119 (1993).
- H. Mooney and G. Koch, *Ambio* 23, 74 (1994); B. Walker, *ibid.*, p. 67.
 - Only reports presenting direct field observations were included in the data, and no conversion factors were used to recalculate values. Detrital C mass (grams of C per square meter) was measured as dead plant material for macrophytes, sampled from standard procedures (22), or as dead cells for microalgae (22). Production of detrital C (C flux to the detrital pool, in grams of C per square meter per day) was calculated as the sum of either litterfall for macrophytes or nongrazing cell mortality for microalgae (23), and plant exudation of dissolved organic C (7). Areal values of phytoplanktonic C flux into the detrital pool were obtained by integration over the mixed layer. Detrital C turnover rate (per day), which at a steady state indicates the fractional loss rate of C from the detrital pool (8), was calculated as the ratio of C flux into the detrital pool to detrital C mass. Plant turnover rate (primary production per unit of plant biomass, per day) was used as an estimate of relative growth rate (24). We excluded short-term studies (that is, studies shorter than the growth season of the dominant primary producer), because they could not be extrapolated to an annual basis, and studies focused on single species, unless that species dominated ecosystem biomass and primary production.
 - O. Vadstein, B. Harkjær, A. Jensen, Y. Olsen, H. Reinertsen, *Limnol. Oceanogr.* 34, 840 (1989); K. Khalilov and Z. Buriakova, *ibid.* 14, 521 (1969); R. Wetzel and P. Penhale, *Aquat. Bot.* 6, 149 (1979); C. Grier and R. Logan, *Ecol. Monogr.* 47, 373 (1977).
 - J. Proctor, J. Anderson, P. Chai, H. Vallack, *J. Ecol.* 71, 237 (1983); M. Piller and J. Stuart, *Can. J. For. Res.* 23, 552 (1993).
 - R. Van Hook, *Ecol. Monogr.* 41, 1 (1971); S. Cargill and R. Jefferies, *J. Appl. Ecol.* 21, 669 (1984); S. Lee, *Mar. Ecol. Prog. Ser.* 66, 161 (1990); M. Lowman, *J. Ecol.* 76, 45 (1988); W. Harris, P. Sollins, N. Edwards, B. Dinger, H. Shugart, in *Productivity of the World Systems. Proceedings of a Symposium*, D. Reichle, J. Franklin, D. Goodall, Eds. (National Academy of Sciences, Washington, DC, 1975), pp. 116–122; L. Bliss, in *Structure and Function of Tundra Ecosystems*, T. Rosswall and O. Heal, Eds. (Tundra Biome Steering Committee, Stockholm, 1975), pp. 17–60; R. Wiegert and F. Evans, *Ecology* 45, 49 (1964); S. McNaughton, *Ecol. Monogr.* 55, 259 (1985); R. Strugnell and C. Pigott, *J. Ecol.* 66, 73 (1978); E. Menhinick, *Ecol. Monogr.* 37, 255 (1967); D. Coppock, J. Dettling, M. Dyer, *Oecologia (Berlin)* 66, 1 (1983); A. McLachlan and G. Bate, *Vie Milieu* 34, 67 (1985); J. Magonigal and F. Day, *Am. J. Bot.* 75, 1334 (1988); J. Frangi and A. Lugo, *Ecol. Monogr.* 55, 351 (1985); F. Schalles and L. Shure, *ibid.* 59, 365 (1989); G. Pellikaan and P. Nienhuis, *Aquat. Bot.* 30, 189 (1988); R. Ratliff and S. Westfall, *ibid.*, p. 109; W. Granell, *ibid.* 35, 99 (1989); J. Van Wijk, *ibid.* 31, 211 (1988); R. Lignell et al., *Mar. Ecol. Prog. Ser.* 94, 239 (1993); B. Wheeler and S. Shaw, *J. Ecol.* 79, 285 (1991); J. Karagatzides and I. Hutchinson, *ibid.*, p. 459; J. Romero, G. Pergent, C. Pergent-Martini, M. Mateo, C. Regnier, *Mar. Ecol.* 13, 69 (1992).
 - G. Cox and J. Whaitaka, *Oikos* 54, 60 (1989).
 - M. Brylinsky, *Mar. Biol.* 39, 213 (1977); P. Penhale and W. Smith, *Limnol. Oceanogr.* 22, 400 (1977); M. Covehey, M. Enell, K. Larsson, L. Olofsson, *Oikos* 29, 5 (1977); J. Proctor, J. Anderson, S. Fogden, H. Vallack, *J. Ecol.* 71, 261 (1983); D. Madge, *Pedobiologia* 5, 273 (1965); R. Chew and A. Chew, *Ecol. Monogr.* 35, 355 (1965); *ibid.* 40, 1 (1970); R. Douglas-Hunter, *Hydrobiologia* 69, 251 (1980); E. Tanner, *J. Ecol.* 68, 833 (1980); D. Reichle and A. Crossley, in *Secondary Productivity of Terrestrial Ecosystems*, K. Petrusewicz, Ed. (Institute of Ecology, Polish Academy of Sciences, Warszawa, 1967), pp. 563–581; C. Mason and R. Bryant, *Freshwater Biol.* 5, 271 (1975); R. Carpenter, *Ecol. Monogr.* 56, 345 (1986); A. Jespersen, K. Christoffersen, B. Riemann, *Verh. Int. Verein. Theor. Angew. Limnol.* 23, 440 (1988); S. Sheldon, *Ecology* 68, 1920 (1987); A. Sinclair, *J. Anim. Ecol.* 44, 497 (1975); W. Darley, C. Montague, F. Plumley, W. Sage, A. Psalidas, *J. Phycol.* 17, 122 (1981); R. Gulati, *Verh. Int. Verein. Theor. Angew. Limnol.* 19, 1202 (1975); Z. Gliwicz and A. Hillbricht-Ilkowska, *ibid.* 18, 197 (1972); D. Cushing, *Limnol. Oceanogr.* 21, 359 (1976); J. Klopatek and F. Stearns, *Am. Midl. Nat.* 100, 320 (1978); O. Heal, in *Tundra Biome. Proceedings IV International Meeting on the Biological Productivity of Tundra*, Swedish International Biological Programme Committee, Leningrad, October 1971, F. Wielgolaski and T. Rosswall, Eds. (Tundra Biome Steering Committee, Stockholm, 1972), pp. 295–305; W. Lampert, *Verh. Int. Verein. Theor. Angew. Limnol.* 20, 969 (1978); J. Gray, *Ecol. Monogr.* 52, 415 (1982); M. Nakanishi et al., *Arch. Hydrobiol. Beih. Ergeb. Limnol.* 35, 47 (1992); B. Jupp and D. Spence, *J. Ecol.* 65, 431 (1977); R. Whittaker, *Communities and Ecosystems* (Collier Macmillan, London, 1975), p. 240; J. Jacoby, *Freshwater Biol.* 18, 495 (1987); M. Nicotri, *Ecology* 58, 1020 (1977); H. Skjoldal and C. Lännergren, *Mar. Biol.* 47, 313 (1978); G. Soszka, *Ecol. Pol.* 23, 371 (1975); D. Baird and R. Ulanowicz, *Ecol. Monogr.* 59, 329 (1989); J. Wanders, *Aquat. Bot.* 3, 357 (1977); K. Mann, *Mar. Biol.* 14, 199 (1972); D. Klumpp, J. Salita-Espinosa, M. Fortes, *Aquat. Bot.* 43, 327 (1992); M. Gandar, *Oecologia (Berlin)* 54, 370 (1982); J. Möller and S. Schwaerter, *Verh. Int. Verein. Theor. Angew. Limnol.* 23, 445 (1988); A. Cattaneo, *Limnol. Oceanogr.* 28, 124 (1983); P. Nienhuis and A. Groenendijk, *Mar. Ecol. Prog. Ser.* 29, 29 (1986); J. Ott and L. Maurer, in *Biology of Benthic Organisms*, B. Keegan, Ed. (Pergamon, Oxford, 1977), pp. 493–502; D. Klumpp, R. Howard, D. Pollard, in *Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region*, A. Larkum, A. McComb, S. Sæpheard, Eds. (Elsevier, Amsterdam, 1989), pp. 394–457; J. Bray, *Ecology* 45, 165 (1964); J. Teal, *ibid.* 43, 614 (1962); M. Tilzer, *J. Plankton Res.* 6, 309 (1984); D. Kirkman and D. Reid, *Aquat. Bot.* 7, 173 (1979); M. Anderson and J. Low, *J. Wildl. Manage.* 40, 233 (1976); J. Ogden, in *Handbook of Seagrass Biology: An Ecosystem Approach*, R. Phillips and C. McRoy, Eds. (Garland, New York, 1980), pp. 173–198; R. Newell, J. Field, C. Griffiths, *Mar. Ecol. Prog. Ser.* 8, 103 (1982); R. Jacobs, C. Den Hartog, B. Braster, F. Carriere, *Aquat. Bot.* 10, 241 (1981); D. Stanley, *Ecology* 57, 1015 (1976); D. Stanley, *ibid.*, p. 1034; R. Flint and C. Goldman, *Limnol. Oceanogr.* 20, 935 (1975); W. Admiraal, L. Bouwman, L. Hoekstra, K. Romeyn, *Int. Rev. Gesamten Hydrobiol.* 68, 175 (1983); J. Garnier and S. Mourelatos, *Freshwater Biol.* 25, 515 (1991); D. Jewson, B. Rippey, W. Gilmore, *Limnol. Oceanogr.* 26, 1045 (1981); J. Wallace and J. O'Hop, *Ecology* 66, 1534 (1985); M. Greenway, *Aquat. Bot.* 2, 117 (1976); D. Cahoon and R. Stevenson, *J. Ecol.* 67, 1341 (1986); C. Ohmart, L. Stewart, J. Thomas, *Oecologia (Berlin)* 59, 322 (1983); B. Frost, *Limnol. Oceanogr.* 36, 1616 (1991); E. Laws, P. Bienfang, D. Ziemann, L. Conquest, *ibid.* 33, 57 (1988); T. Weiss et al., *ibid.* 35, 781 (1990); J. Cloern, B. Cole, S. Wienke, *ibid.* 32, 815 (1987); N. Welschmeyer and C. Lorenzen, *ibid.* 30, 1 (1985); J. Macanley, J. Clark, A. Price, *Aquat. Bot.* 31, 277 (1988); H. Odum, *Ecol. Monogr.* 27, 55 (1957); E. Leigh and D. Windsor, in *Ecology of a Tropical Forest: Seasonal Rhythms and Long Term Changes*, E. Leigh, A. Raud, D. Windsor, Eds. (Smithsonian Institution Press, Washington, DC, 1983), pp. 111–123; D. Klumpp, J. Salita-Espinosa, M. Fortes, *Aquat. Bot.* 45, 205 (1993).
 - The relation between C flux into the detrital pool (FC_{detrital} , grams of C per square meter per day) and primary production (PP, grams of C per square meter per day) for all the ecosystem types is described by the equation

$$\log FC_{\text{detrital}} = (-0.25 \pm 0.04) + (1.00 \pm 0.04) \log PP$$
 with r^2 (r is the Pearson correlation coefficient) = 0.84, F test, $P < 0.01$, and $n = 121$. The slopes ($\pm SE$) of the log-log linear regression equations be-
 - tween C flux into the detrital C pool and primary production fitted to each ecosystem type are as follows: plankton, 1.54 ± 0.25 ($P < 0.0001$, $r^2 = 0.61$, $n = 24$); benthic microalgae, 1.12 ± 0.07 ($P < 0.0001$, $r^2 = 0.96$, $n = 16$); macroalgae beds, 0.97 ± 0.03 ($P < 0.0001$, $r^2 = 0.99$, $n = 13$); freshwater macrophyte meadows, 1.19 ± 0.09 ($P < 0.0001$, $r^2 = 0.93$, $n = 16$); seagrass meadows, 0.99 ± 0.05 ($P < 0.0001$, $r^2 = 0.99$, $n = 6$); grasslands, 0.50 ± 0.016 ($P < 0.01$, $r^2 = 0.42$, $n = 13$); and forests 0.98 ± 0.07 ($P < 0.0001$, $r^2 = 0.99$, $n = 33$). All of the slopes were not significantly different from 1 (null hypothesis, H_0 : slope = 1, t test, $P > 0.1$), except the slope for grasslands. The relation between C flux into the detrital pool and primary production for grasslands is influenced by two reports from the African savanna (70), where herbivory is disproportionately higher than in the remaining grassland systems. These high levels of herbivory greatly reduce detrital C flux resulting in a nonlinear relation between C flux into the detrital pool and primary production for the grasslands in the data.
 - D. DeAngelis, R. Gardner, H. Shugart, in *Dynamics and Properties of Forest Ecosystems*, D. Reichle, Ed. (Cambridge Univ. Press, Cambridge), pp. 567–672.
 - M. Silver and M. Gowing, *Prog. Oceanogr.* 26, 75 (1991); V. Asper, W. Deuser, G. Knauer, S. Lohrenz, *Nature* 357, 670 (1992); R. Lampitt, *Deep-Sea Res.* 32, 885 (1985).
 - S. Enriquez, C. Duarte, K. Sand-Jensen, *Oecologia* 94, 457 (1993); P. Coley, J. Bryant, F. Stuart Chapin III, *Science* 230, 895 (1985); F. Stuart Chapin III, A. Bloom, C. Field, R. Waring, *BioScience* 37, 49 (1987).
 - K. Sand-Jensen and J. Borum, *Aquat. Bot.* 41, 137 (1991); J. Neundorfer and W. Kemp, *Mar. Ecol. Prog. Ser.* 94, 71 (1993); C. Duarte, *Ophelia* 41, 87 (1995).
 - S. Subak, P. Raskin, D. Von Hippel, *National Greenhouse Gas Accounts: Current Anthropogenic Sources and Sinks* (Stockholm Environment Institute, Stockholm, Sweden, 1992), p. 37.
 - J. Overpeck, P. Bartlein, T. Webb III, *Science* 254, 692 (1991).
 - O. Phillips and A. Gentry, *ibid.* 263, 954 (1994).
 - H. Polley, B. Johnson, H. Marino, S. Mayeux, *Nature* 361, 61 (1993); J. Van der Burgh, H. Visscher, D. Dilcher, W. Kürschner, *Science* 260, 1788 (1993).
 - F. Bazzaz, *Annu. Rev. Ecol. Syst.* 21, 167 (1990); H. Mooney, B. Drake, R. Luxmoore, W. Oechel, L. Pitelka, *BioScience* 41, 96 (1991); C. Körner and J. Arnone III, *Science* 257, 1672 (1992).
 - Dead plant material was calculated as the sum of standing dead material, still attached to the plant, and shed material on the floor. Most reports quantified the mass of dead plant material on the floor after samples collected from the soil surface were sieved through a mesh with a pore size of 1 mm. This procedure underestimates detrital C mass in the soil because it ignores plant material <1 mm and that buried below the soil surface. The detrital mass of planktonic and benthic microalgae represents particulate detritus alone, because it is measured as the abundance of dead cells.
 - Nongrazing cell mortality was quantified as the sum of cell sedimentation rates and lysis rates. Direct estimates of cell lysis are very scarce, and most of the compiled reports on microalgal lysis rates were derived from the following carbon budget equation: primary production = biomass increment + grazed production + exuded production + sedimentation rate + exported production by advection + lysis rate (all variables in grams of C per square meter per day).
 - J. Cebrían and C. M. Duarte, *Funct. Ecol.* 8, 518 (1994).
 - We thank K. Sand-Jensen, J. Kalff, R. Peters, J. Kenworthy, Y. Prairie, and T. Granata for improving the manuscript. This work was supported by the Spanish Interministerial Commission of Science and Technology (CICYT) and a fellowship from the Comissió Interdepartamental de Recerca i Innovació Tecnològica to J.C.